10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-Cat IgG F(ab')2 (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin
Catalog # ASR2149

## Product Information

| | |
|---|---|
| **Description** | Anti-CAT IgG F(ab')2 (GOAT) Antibody Biotin Conjugated |
| **Host** | Goat |
| **Conjugate** | Biotin |
| **Target Species** | Cat |
| **Clonality** | Polyclonal |
| **Physical State** | Lyophilized |
| **Host Isotype** | IgG |
| **Target Isotype** | IgG F(ab')2 |
| **Buffer** | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| **Immunogen** | Cat IgG F(ab')2 fragment |
| **Reconstitution Volume** | 1.0 mL |
| **Reconstitution Buffer** | Restore with deionized water (or equivalent) |
| **Stabilizer** | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| **Preservative** | 0.01% (w/v) Sodium Azide |

## Additional Information

| | |
|---|---|
| **Shipping Condition** | Ambient |
| **Application Note** | This product has been assayed against 1.0 ug of Cat IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 is suggested for this product. |
| **Purity** | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Cat IgG, Cat IgG F(ab')2 and Cat Serum. No reaction was observed against Cat IgG F(c). |
| **Storage Condition** | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| **Precautions Note** | This product is for research use only and is not intended for therapeutic or diagnostic applications. |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.