10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



F(ab')2 Anti-Mouse IgG F(c) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2247

Product Information

Description F(ab')2 Anti-MOUSE IgG F(c) (GOAT) Antibody Biotin Conjugated Min X

Bv Hs & Hu Serum Proteins

HostGoatConjugateBiotinTarget SpeciesMouseClonalityPolyclonalPhysical StateLyophilizedHost IsotypeIgG F(ab')2

Target Isotype IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Mouse IgG F(c) fragment

Reconstitution Volume 500 □

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Application Note This product has been assayed against 1.0 ug of Mouse IgG in a standard

capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.

Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(ab')2 or Bovine, Horse and

Human Serum Proteins.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.