

Horse IgG Fab Rhodamine

Catalog # ASR2262

Product Information

Description HORSE IgG F(ab) fragment Rhodamine conjugated

Conjugate Rhodamine (TRITC)

FP Value 1.0 moles Rhodamine (TRITC) per mole of Horse IgG F(ab)

Physical StateLyophilizedHost IsotypeIgG F(ab)

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Species of Origin Horse **Reconstitution Volume** 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Purity This product was prepared from normal serum by delipidation, salt

fractionation, ion exchange chromatography followed by papain digestion

and extensive dialysis against the buffer stated above. Assay by

immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse

IgG, anti-Mouse IgG F(ab')2 and anti-Mouse Serum. No reaction was

observed against anti-Mouse IgG F(c) or anti-Pepsin.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.