

F(ab')2 Anti-Cat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody

Rabbit Polyclonal, Peroxidase (Horseradish) Catalog # ASR2376

Product Information

Description F(ab')2 Anti-CAT IgG [H&L] (RABBIT) Antibody Peroxidase Conjugated

Host Rabbit

Conjugate Peroxidase (Horseradish)

Target SpeciesCatReactivityCatClonalityPolyclonalPhysical StateLyophilized

Host Isotype IgG F(ab')2

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Anti-Cat IgG (H&L) was produced by repeated immunization with Cat IgG

whole molecule in rabbit.

Reconstitution Volume 500 □

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Additional Information

Shipping Condition Ambient

Application Note Antibody anti-Cat has been assayed against 1.0 ug of Cat IgG in a standard

capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product. Antibody anti-Cat has been assayed against 1.0 ug

of Cat IgG in a standard capture ELISA using ABTS

(2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product.

Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Cat IgG and Cat Serum. No reaction was observed against anti-Pepsin or anti-Rabbit

IgG F(c).

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and

thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

Anti-Rat IgG (H&L) antibody generated in rabbit detects specifically Cat IgG (H&L). This secondary antibody anti-Cat is ideal for investigators who routinely perform titration assays, westCrn-blot, immunoprecipitation and more generally immunoassays.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.