10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish) Catalog # ASR2389

Product Information

Description F(ab')2 Anti-HUMAN IgG F(ab')2 (GOAT) Antibody Peroxidase

Conjugated Min X Bv Hs Ms & Rt Serum Proteins

Host Goat

Conjugate Peroxidase (Horseradish)

Target Species Human Reactivity Human Clonality Polyclonal **Physical State** Lyophilized IgG F(ab')2 **Host Isotype**

Target Isotype IgG F(ab')2

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Anti-Human IgG F(ab')2 fragment antibody was produced by repeated **Immunogen**

immunization with Human IgG F(ab')2 fragment in goat.

Reconstitution Volume 500 □

Reconstitution Buffer

Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Additional Information

Shipping Condition Ambient

Suitable for immunoblotting (western or dot blot), ELISA, immunoelectron **Application Note**

microscopy and immunohistochemistry as well as other antibody based

enzymatic assays requiring lot-to-lot consistency.

This product was prepared from monospecific antiserum by immunoaffinity Purity

> chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin

digestion and chromatographic separation. Assay by

immunoelectrophoresis resulted in a single precipitin arc against

anti-Peroxidase, anti-Goat Serum, Human IgG, Human IgG F(ab')2 and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c),

Human IgG F(c) or Bovine, Horse, Mouse and Rat Serum Proteins.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

> contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

Anti-Human IgG F(ab')2 fragment antibody generated in goat detects specifically Human IgG F(ab')2 fragment. This secondary antibody anti-Human is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.