

F(ab')₂ Anti-Mouse IgG F(c) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish)

Catalog # ASR2524

Product Information

Description	F(ab') ₂ Anti-MOUSE IgG F(c) (GOAT) Antibody Peroxidase Conjugated Min X Bv Hs & Hu Serum Proteins
Host	Goat
Conjugate	Peroxidase (Horseradish)
Target Species	Mouse
Clonality	Polyclonal
Physical State	Lyophilized
Host Isotype	IgG F(ab') ₂
Target Isotype	IgG F(c)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Mouse IgG F(c) fragment
Reconstitution Volume	500 μ L
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Additional Information

Shipping Condition	Ambient
Application Note	F(ab') ₂ Anti-Mouse IgG F(c) Antibody is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 ⁶ cells in flow cytometry is approximately 1.0 μ g of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.
Purity	F(ab') ₂ Anti-Mouse IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(ab') ₂ or Bovine, Horse, and Human Serum Proteins.
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

F(ab')₂ Anti-Mouse IgG F(c) Antibody generated in goat specifically detects Mouse IgG F(ab')₂. F(ab')₂ Anti-Mouse IgG F(c)Antibody is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. F(ab')₂ Anti-Mouse IgG F(c) Antibody is ideal for investigators in Cancer, Immunology, and Microbiology research.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.