

Anti-Mouse IgG3 (Gamma 3 chain) (Texas Red™ Conjugated) Secondary Antibody

Rabbit Polyclonal, Texas Red® Catalog # ASR2856

Product Information

Description Anti-MOUSE IgG3 (Gamma 3 chain) (RABBIT) Antibody Texas Red™

Conjugated

HostRabbitConjugateTexas Red®

FP Value 3.0 moles Texas Red® per mole of IgG

Target SpeciesMouseClonalityPolyclonalPhysical StateLyophilized

Host Isotype IgG
Target Isotype IgG3

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Mouse IgG3 heavy chain

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Mouse antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum and Mouse Serum. Specificity was confirmed by ELISA at less than 1% cross reactivity against other mouse heavy or light chain isotypes.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and

fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.