

F(ab')₂ Anti-Rabbit IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody

Goat Polyclonal, R-Phycoerythrin (RPE)

Catalog # ASR2871


Product Information

Description	F(ab') ₂ Anti-RABBIT IgG [H&L] (GOAT) Antibody Phycoerythrin conjugated
Host	Goat
Conjugate	R-Phycoerythrin (RPE)
Target Species	Rabbit
Clonality	Polyclonal
Physical State	Lyophilized
Host Isotype	IgG F(ab') ₂
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Rabbit IgG whole molecule
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition	Ambient
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Coupling to R-PE was followed by size exclusion chromatography to purify conjugate from unreacted R-PE and antibody. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Goat Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).
Storage Condition	Store vial at 4° C prior to opening. Dilute only prior to immediate use. Do not freeze after reconstitution. Store reagent in the dark. This product is stable at 4° C as an undiluted liquid. Use subdued lighting during handling and incubation of cells prior to analysis.
Precautions Note	This product is for research use only and is not intended for therapeutic or diagnostic applications.

Images



Dot Blot of F(ab)2 Goat anti-Rabbit IgG Antibody
Phycoerythrin Conjugated. Antigen: Row 1: Rabbit IgG
Row 2: Goat IgG Row 3: Mouse IgG Load: Lane 1 - 200ng
Lane 2 - 66.7ng Lane 3 - 22.2ng Lane 4 - 7.4ng Lane 5 -
2.5ng Primary antibody: N/A Secondary antibody: F(ab)2
Goat anti-Rabbit IgG Antibody Phycoerythrin Conjugated
at 1:1,000 for 60 min at RT. Block: MB-070 for 1 hour at
RT

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.