

Chicken IgG F(c)

Catalog # ASR2895

Product Information

| Description Conjugate Physical State Host Isotype Buffer Species of Origin Stabilizer Preservative | CHICKEN IgG F(c) fragment Unconjugated Lyophilized IgG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Chicken None |
|---|---|
| Preservative | 0.01% (w/v) Sodium Azide |

Additional Information

| Shipping Condition | Wet Ice |
|--------------------|---|
| Purity | This product was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and papain digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Chicken Serum, anti-Chicken IgG and anti-Chicken IgG F(c). No reaction was observed against anti-Chicken IgG F(ab')2 or anti-Papain. |
| Storage Condition | Store vial at 4° C prior to restoration. Restore with 1.0 mL of deionized water (or equivalent). For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Precautions Note | This product is for research use only and is not intended for therapeutic or diagnostic applications. |

Images



SDS-PAGE of Chicken IgG F(c) Fragment (p/n ASR2895). Lane M: 5 µL Opal Prestained Marker (p/n MB-210-0500). Lane 1: Reduced Chicken IgG Whole Molecule (p/n 003-0102). Lane 2: Reduced Chicken IgG F(c) Fragment (p/n ASR2895). Lane 3: Reduced Chicken IgG F(ab) Fragment (p/n 003-0105). Lane 4: Reduced Chicken IgM Whole Molecule (p/n 003-0107). Load: 1 µg per lane. Predicted/Observed size: IgG at 72 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 75 kDa. Observed F(c) Fragment migrates slightly higher. Other bands: Chicken IgG heavy chain alternative splicing variant at approximately 40 kDa in Lane 1.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.