

Guinea Pig IgM

Catalog # ASR2897

Product Information

| | |
|--------------------------|--|
| Description | GUINEA PIG IgM whole molecule |
| Conjugate | Unconjugated |
| Physical State | Liquid (sterile filtered) |
| Host Isotype | IgM |
| Buffer | 0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0 |
| Species of Origin | Guinea Pig |
| Preservative | 0.1% (w/v) Sodium Azide |


Additional Information

| | |
|---------------------------|---|
| Shipping Condition | Wet Ice |
| Application Note | Guinea Pig IgM whole molecule can be utilized as a control or standard reagent in Western Blotting and ELISA experiments. |
| Purity | Guinea Pig Igm whole molecule was prepared from normal serum by a multi-step process which includes delipidation, selective precipitation and tandem molecular sieve chromatography followed by extensive dialysis against the buffer stated above. Guinea Pig Igm whole molecule was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Guinea Pig Serum and anti-Guinea Pig IgM (λ chain specific). No reaction was observed against anti-Guinea Pig IgG F(c). Some light chain cross reactivity will occur with anti-Guinea Pig IgG. |
| Storage Condition | Store vial at 4° C prior to opening. Guinea Pig Igm whole molecule is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. |
| Precautions Note | This product is for research use only and is not intended for therapeutic or diagnostic applications. |

Background

Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum.

Images



SDS-Page of Guinea Pig IgM Whole Molecule. Lane 1: Guinea Pig IgM, Non-Reduced. Lane 2: Guinea Pig IgM, Reduced. Load: 1.0 μ g per lane. Predicted/Observed size-Predicted/Observed size - Non-Reduced: 900 kDa (Pentamer), 900 kDa (Molecule larger than can pass through gel), Reduced: 78 and 25 kDa, 75 and 25 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.