10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



## Anti-Goat IgM (mu) (Peroxidase Conjugated) Secondary Antibody

Rabbit Polyclonal, Peroxidase (Horseradish) Catalog # ASR2914

## **Product Information**

**Description** Anti-GOAT IgM (mu) (RABBIT) Antibody Peroxidase Conjugated

**Host** Rabbit

**Conjugate** Peroxidase (Horseradish)

Target SpeciesGoatClonalityPolyclonalPhysical StateLyophilized

Host Isotype IgG

Target Isotype IgM □chain

**Buffer** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Immunogen** Goat IgM mu heavy chain

**Reconstitution Volume** 2.0 mL

**Reconstitution Buffer** Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

**Preservative** 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

## Additional Information

Shipping Condition Ambient

**Application Note** This product has been assayed against 1.0 ug of Goat IgM in a standard

capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:7,000 to 1:35,000 of the reconstitution concentration is

suggested for this product.

**Purity** This product is an IgG fraction antibody purified from monospecific

antiserum by a multi-step process which includes delipidation, salt

fractionation and ion exchange chromatography followed by cross adsorption to remove any unwanted reactivities and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Goat IgM and Goat Serum.

No reaction was observed against Goat IgG.

**Storage Condition** Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

**Precautions Note**This product is for research use only and is not intended for therapeutic or

diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.