10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



Anti-Golden Syrian Hamster IgM mu chain (Peroxidase Conjugated) Secondary Antibody

Rabbit Polyclonal, Peroxidase (Horseradish) Catalog # ASR2917

Product Information

Description Anti-GOLDEN SYRIAN HAMSTER IgM mu chain (RABBIT) Antibody Peroxidase

Conjugated

Host Rabbit

ConjugatePeroxidase (Horseradish)Target SpeciesGolden Syrian Hamster

Clonality Polyclonal **Physical State** Lyophilized

Host Isotype IgG

Target Isotype IgM □chain

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Hamster IgM mu heavy chain

Reconstitution Volume 2.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Additional Information

Shipping Condition Ambient

Application Note This product has been assayed against 1.0 ug of Hamster IgM in a standard

capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:12,000 of the reconstitution concentration is

suggested for this product.

Purity This product is an IgG fraction antibody purified from monospecific

antiserum by a multi-step process which includes delipidation, salt

fractionation and ion exchange chromatography followed by cross adsorption to remove any unwanted reactivities and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Hamster IgM and Hamster

Serum. No reaction was observed against Hamster IgG.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.