10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999

abcepta **

Anti-Monkey IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR3007

Product Information

Description Anti-MONKEY IgG IgA IgM (H&L) (GOAT) Antibody Peroxidase Conjugated

Host Goat

Conjugate Peroxidase (Horseradish)

Target SpeciesMonkeyClonalityPolyclonalPhysical StateLyophilized

Host Isotype IgG

Target Isotype IgG IgA IgM

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Anti-Monkey IgG, IgA and IgM were produced by repeated immunization

with monkey IgG, IgA, IgM whole molecules in goat.

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Thimerosal

Additional Information

Shipping Condition Ambient

Application Note Anti-Monkey IgA, IgG and IgM peroxidase conjugated antibody are suitable

for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody

based enzymatic assays requiring lot-to-lot consistency.

Purity This product was prepared from polyspecific antiserum by immunoaffinity

chromatography using antigens coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Monkey IgG, Monkey IgA and Monkey IgM. This reagent is suitable for the detection of all Monkey immunoglobulin

subclasses and chain combinations.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Background

Anti-Monkey IgG, IgA and IgM peroxidase conjugated antibodies generated in goat detect specifically Monkey IgA, IgG and IgM. These secondary peroxidase conjugated antibodies anti-Monkey are ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.

Images

Product	Gt-a-Monkey IgG	Gt-a-Monkey IgA	Gt-a-Monkey IgM
Rhesus Serum	+	+	+
Baboon Serum	+	+	+
Cynomologus Serum	+	+	+
Rhesus IgG	+		
Human IgA		+	
Human IgM			+

This table displays additional reactivity among various species of serum and immunoglobulin. A (+) indicates antibody reactivity to the corresponding target.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.