

F(ab')2 Anti-Rat IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody

Donkey Polyclonal, Peroxidase (Horseradish) Catalog # ASR3175

Product Information

Description	F(ab')2 Anti-RAT IgG [H&L] (DONKEY) Antibody Peroxidase Conjugated Min X Bv Ch Gt GP Ham Hs Hu Ms Rb & Sh Serum Proteins
Host	Donkey
Conjugate	Peroxidase (Horseradish)
Target Species	Rat
Clonality	Polyclonal
Physical State	Lyophilized
Host Isotype	IgG F(ab')2
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Rat IgG whole molecule
Reconstitution Volume	500 L
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Additional Information

Shipping Condition	Ambient
Application Note	Suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Do not use sodium azide as a preservative. Azide will substantially inhibit the enzyme activity of horseradish peroxidase.
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c), or Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Sheep serum proteins.
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.