

Anti-NEDD1 Secondary Antibody

Rabbit Polyclonal, Unconjugated

Catalog # ASR3293

Product Information

| | |
|-----------------------|---|
| Description | Anti-NEDD1 (RABBIT) Antibody |
| Host | Rabbit |
| Conjugate | Unconjugated |
| Target Species | Human |
| Reactivity | Human |
| Clonality | Polyclonal |
| Application | WB |
| Physical State | Liquid (sterile filtered) |
| Host Isotype | Antiserum |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Immunogen | Anti-NEDD1 was prepared from whole rabbit serum produced by repeated immunizations with a recombinant protein corresponding to the 343-667 region of human Nedd1. |
| Stabilizer | None |
| Preservative | 0.01% (w/v) Sodium Azide |


Additional Information

| | |
|---------------------------|--|
| Shipping Condition | Dry Ice |
| Application Note | This antiserum has been tested for use in ELISA and western blotting using a recombinant truncated Nedd1 protein. Specific conditions for reactivity and detection of Nedd1 should be optimized by the end user. Expect a band approximately ~73 kDa in size corresponding to Nedd1 by western blotting in the appropriate cell lysate or extract. |
| Purity | This product was adsorbed against GST from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with endogenous Nedd1 protein. A BLAST analysis was used to suggest reactivity with Nedd1 from human, chimpanzee, macaque, marmoset, cattle, rat, and mouse based on a 100% homology with the immunizing sequence. Expect partial reactivity with Nedd1 from turkey, chicken, salmon, and Danio based on a 91% homology with the immunizing sequence. Cross-reactivity with Nedd1 from other sources has not been determined. |
| Storage Condition | Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Precautions Note | This product is for research use only and is not intended for therapeutic or diagnostic applications. |

Background

This antibody is designed, produced, and validated as part of a collaboration between Rockland and the National Cancer Institute (NCI) and is suitable for Cancer, Immunology and Nuclear Signaling research. Microtubules are polymers of tubulin, which exist as heterodimers of alpha-tubulin and beta-tubulin. NEDD1 (neural precursor expressed, developmentally down-regulated protein1; also called GCP-WD) is a centrosomal protein that in mammals associates with the gamma-tubulin ring complex (γ -TuRC). γ -TuRC is critical for initiation, or nucleation, of the microtubule assembly. In association with this complex, NEDD1 plays an important role in targeting the γ -TuRC complex to the site of microtubule nucleation and to the mitotic spindle. These events are essential for proper bipolar spindle formation and mitotic progression. Given the casual link between improper spindle function and tumorigenesis, characterization of Nedd1 function will be important to better understand various mechanisms underlying mitotic regulation, chromosome segregation, and cancer development.

Images



Anti-NEDD1 in Western Blot using Abcepta Immunochemicals' Anti-NEDD1 Antibody shows detection of a 73 kDa band corresponding to endogenous NEDD1 in lysates of S phase HeLa cells silenced for either control Luciferase or NEDD1. In right lane (NEDD1i): lysates from sh-NEDD1 RNAi-treated lentivirus-infected cells. In left lane (GLi): lysates from sh-Luciferase lentivirus-infected cells as control. Anti-NEDD1 Antibody was used at 1:10,000. Molecular weight estimation was made by comparison by prestained MW markers. ECL was used for detection. Personal communication, Kyung S. Lee, NCI, Bethesda, MD.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.