

Anti-Rat IgG F(ab')2 Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR3334

Product Information

Description Anti-RAT IgG F(ab')2 (RABBIT) Antibody

Host Rabbit

Conjugate Unconjugated

Target SpeciesRatClonalityPolyclonalApplicationWB

Physical State Liquid (sterile filtered)

Host Isotype IgG

Target Isotype IgG F(ab')2

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Rat IgG F(ab')2 fragment

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Wet Ice

Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by

immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit

Serum, Rat IgG, Rat IgG F(ab')2 and Rat Serum. No reaction was observed

against Rat IgG F(c).

Storage Condition Store vial at 4° C prior to opening. This product is stable for several weeks

at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid


cycles of freezing and thawing.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Images

Western Blot of Anti-Rat IgG F(ab')2 (RABBIT) Antibody (p/n ASR3334). Lane M: 3 μ l Molecular Ladder. Lane 1: Rat IgG whole molecule (p/n 012-0102). Lane 2: Rat IgG F(c) Fragment (p/n 012-0103). Lane 3: Rat IgG F(ab) Fragment (p/n 012-0105). Lane 4: Rat IgM Whole Molecule (p/n 012-0107). All samples were reduced. Load: 50 ng per



lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rat IgG F(ab')2 (RABBIT) Antibody (p/n ASR3334) 1:1,000 for 60 min at RT. Secondary Antibody: Anti-Rabbit IgG (GOAT) Peroxidase Conjugated Antibody (p/n 611-103-122) 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 55 kDa for Rat IgG, 25 kDa for F(c) and F(ab), 78 and 25 kDa for IgM. Rat F(c) migrates slightly higher.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.