

Anti-Mouse IgG F(c) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR3534

Product Information

Description Anti-MOUSE IgG F(c) (GOAT) Antibody

Host Goat

Conjugate Unconjugated

Target SpeciesMouseReactivityMouseClonalityPolyclonalPhysical StateLyophilizedHost IsotypeIgG

Host Isotype IgG
Target Isotype IgG F(c)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Anti-Mouse IgG was produced by repeated immunization with mouse IgG

F(c) fragment in goat.

Reconstitution Volume 5.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Application Note Anti-Mouse IgG F(c) antibody has been assayed against 1.0 ug of Mouse IgG

in a standard capture ELISA using ABTS

(2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product.

Purity Anti-Mouse IgG F(c) Antibody is an IgG fraction antibody purified from

monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG

F(ab')2.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Background

Anti-Mouse IgG F(c) antibody generated in goat detects specifically mouse IgG F(c). This Anti-Mouse IgG F(c) Secondary Antibody is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. Anti-Mouse IgG F(c) Antibody is ideal for investigators in Immunology, Cancer, and Microbiology research.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.