

## Bovine IgG (BULK ORDER) Antibody

Catalog # ASR3553

## **Product Information**

**Description**BOVINE IgG whole molecule (BULK ORDER)

ConjugateUnconjugatedPhysical StateLyophilized

Host Isotype IgG

**Buffer** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Species of Origin Reconstitution Volume**Bovine
5.0 mL

**Reconstitution Buffer** Restore with deionized water (or equivalent)

**Preservative** 0.01% (w/v) Sodium Azide

## **Additional Information**

Shipping Condition Ambient

**Application Note** Bovine IgG whole molecule can be utilized as a control or standard reagent

in Western Blotting and ELISA experiments.

**Purity** Bovine IgG whole molecule was prepared from normal serum by a

multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer

stated above. Bovine IgG whole molecule was assayed by

immunoelectrophoresis resulted in a single precipitin arc against anti-Bovine

Serum and anti-Bovine IgG.

**Storage Condition** Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Bovine IgG whole molecule is stable for several weeks at 4° C

as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note** This product is for research use only and is not intended for therapeutic or

diagnostic applications.

## **Background**

Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-afinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.