

Guinea Pig IgG (BULK ORDER)

Catalog # ASR3568

Product Information

Description	GUINEA PIG IgG whole molecule (BULK ORDER)
Conjugate	Unconjugated
Physical State	Lyophilized
Host Isotype	IgG
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Species of Origin	Guinea Pig
Reconstitution Volume	2.5 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Preservative	0.01% (w/v) Sodium Azide


Additional Information

Shipping Condition	Ambient
Application Note	Guinea Pig IgG whole molecule can be utilized as a control or standard reagent in Western Blotting and ELISA experiments.
Purity	Guinea Pig IgG was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Guinea Pig IgG whole molecule was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Guinea Pig IgG and anti-Guinea Pig Serum.
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Guinea Pig IgG whole molecule is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Precautions Note	This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-afinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present.

Images



SDS-Page of Guinea Pig IgG whole molecule. Lane 1: Guinea Pig IgG – Non-reduced. Lane 2: Guinea Pig IgG – Reduced. Load: 1.0 µg per lane. Predicted/Observed size: 25 & 55 kDa – Reduced, 160 kDa – Non-reduced for IgG whole molecule. Other band(s): None.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.