

ATN1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant ATN1.

Catalog # AT1225a

Product Information

Application	WB, IF, E
Primary Accession	P54259
Other Accession	BC051795
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 Kappa
Clone Names	2C10
Calculated MW	125414

Additional Information

Gene ID	1822
Other Names	Atrophin-1, Dentatorubral-pallidoluysian atrophy protein, ATN1, D12S755E, DRPLA
Target/Specificity	ATN1 (AAH51795, 1 a.a. ~ 110 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IF~~1:50~200 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	ATN1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

Dentatorubral pallidoluysian atrophy (DRPLA) is a rare neurodegenerative disorder characterized by cerebellar ataxia, myoclonic epilepsy, choreoathetosis, and dementia. The disorder is related to the expansion from 7-23 copies to 49-75 copies of a trinucleotide repeat (CAG/CAA) within this gene. The encoded protein includes a serine repeat and a region of alternating acidic and basic amino acids, as well as the variable glutamine repeat. Alternative splicing results in two transcripts variants that encode the same protein.


References

Dentatorubral pallidoluysian atrophy in a Turkish family. Yi? U, et al. Turk J Pediatr, 2009 Nov-Dec. PMID 20196398. Screening for premutation in the FMR1 gene in male patients suspected of spinocerebellar ataxia. Rajkiewicz M, et al. Neurol Neurochir Pol, 2008 Nov-Dec. PMID 19235102. Severe neurological phenotypes of Q129 DRPLA transgenic mice serendipitously created by en masse expansion of CAG repeats in Q76 DRPLA mice. Sato T, et al. Hum Mol Genet, 2009 Feb 15. PMID 19039037. Searching for mutation in the JPH3, ATN1 and TBP genes in Polish patients suspected of Huntington's disease and without mutation in the IT15 gene. Su?ek-Piatkowska A, et al. Neurol Neurochir Pol, 2008 May-Jun. PMID 18651325. Molecular analysis of CAG repeats at five different spinocerebellar ataxia loci: correlation and alternative explanations for disease pathogenesis. Alluri RV, et al. Mol Cells, 2007 Dec 31. PMID 18182848.


Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .



Immunofluorescence of monoclonal antibody to ATN1 on HeLa cell . [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged ATN1 is approximately 0.3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.