

CHST11 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant CHST11. Catalog # AT1530a

Product Information

Application WB, E **Primary Accession** Q9NPF2 Other Accession NM 018413 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 Kappa **Clone Names** 4F1

Clone Names 4F1 Calculated MW 41555

Additional Information

Gene ID 50515

Other Names Carbohydrate sulfotransferase 11, Chondroitin 4-O-sulfotransferase 1,

Chondroitin 4-sulfotransferase 1, C4S-1, C4ST-1, C4ST1, CHST11

Target/Specificity CHST11 (NP_060883, 230 a.a. ~ 337 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions CHST11 Antibody (monoclonal) (M01) is for research use only and not for use

in diagnostic or therapeutic procedures.

Background

Chondroitin 4-sulfotransferases, such as CHST11, catalyze the transfer of sulfate from 3-prime-phosphoadenosine 5-prime-phosphosulfate to position 4 of N-acetylgalactosamine residues in chondroitin (Yamauchi et al., 2000 [PubMed 10722746]).

References

Natural frequency of polymorphisms linked to the chondroitin 4-sulfotransferase genes and its association with placental malaria. Valente B, et al. Trans R Soc Trop Med Hyg, 2010 Oct. PMID 20580388.Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose


JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Inherited genetic variant predisposes to aggressive but not indolent prostate cancer. Xu J, et al. Proc Natl Acad Sci U S A, 2010 Feb 2. PMID 20080650.Identification and characterization of TGFbeta-dependent and -independent cis-regulatory modules in the C4ST-1/CHST11 locus. Willis CM, et al. Genet Mol Res, 2009 Nov 3. PMID 19937589. Coeliac disease-associated risk variants in TNFAIP3 and REL implicate altered NF-kappaB signalling. Trynka G, et al. Gut, 2009 Aug. PMID 19240061.

Images

1.6

1.4 1.2 0.8 0.6 0.4 0.2

0.01



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.62 KDa).

CHST11 monoclonal antibody (M01), clone 4F1 Western Blot analysis of CHST11 expression in K-562 ((Cat # AT1530a)

Western Blot analysis of CHST11 expression in transfected 293T cell line by CHST11 monoclonal antibody (M01), clone 4F1.

Lane 1: CHST11 transfected lysate (Predicted MW: 41.6 KDa).

Lane 2: Non-transfected lysate.

Detection limit for recombinant GST tagged CHST11 is approximately 0.1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.