

CTAG2 Antibody (monoclonal) (M04)

Mouse monoclonal antibody raised against a partial recombinant CTAG2. Catalog # AT1664a

Product Information

ApplicationWB, EPrimary AccessionQ75638Other AccessionNM_020994ReactivityHumanHostMouseClonalitymonoclonalIsotypeIgG2b Kappa

Clone Names 3H1 Calculated MW 21090

Additional Information

Gene ID 30848

Other Names Cancer/testis antigen 2, CT2, Autoimmunogenic cancer/testis antigen

NY-ESO-2, Cancer/testis antigen 62, CT62, L antigen family member 1, LAGE-1,

CTAG2, ESO2, LAGE1

Target/Specificity CTAG2 (NP_066274, 111 a.a. ~ 210 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions CTAG2 Antibody (monoclonal) (M04) is for research use only and not for use

in diagnostic or therapeutic procedures.


Background

This gene encodes an autoimmunogenic tumor antigen that belongs to the ESO/LAGE family of cancer-testis antigens. This protein is expressed in a wide array of cancers including melanoma, breast cancer, bladder cancer and prostate cancer. This protein is also expressed in normal testis tissue. An alternative open reading frame product of this gene has been described in PMID 10399963. This alternate protein, termed CAMEL, is a tumor antigen that is recognized by melanoma-specific cytotoxic T-lymphocytes. Alternate splicing results in multiple transcript variants.


References

Comparisons for detecting NY-ESO-1 mRNA expression levels in hepatocellular carcinoma tissues. Wang XY, et al. Oncol Rep, 2009 Mar. PMID 19212631.Prognostic impact of cancer/testis antigen expression in advanced stage multiple myeloma patients. Andrade VC, et al. Cancer Immun, 2008 Feb 1. PMID 18237105.Identification and expression analysis of novel LAGE-1 alleles with single nucleotide polymorphisms in cancer patients. Shao Y, et al. J Cancer Res Clin Oncol, 2008 Apr. PMID 17899192.A new specific gene expression in squamous cell carcinoma of the esophagus detected using representational difference analysis and cDNA microarray. Kan T, et al. Oncology, 2006. PMID 16446548.The LIFEdb database in 2006. Mehrle A, et al. Nucleic Acids Res, 2006 Jan 1. PMID 16381901.

Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (37.11 KDa) .



Detection limit for recombinant GST tagged CTAG2 is approximately 0.1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.