

CYP7B1 Antibody (monoclonal) (M06)

Mouse monoclonal antibody raised against a partial recombinant CYP7B1.

Catalog # AT1706a

Product Information

Application	WB, IHC, E
Primary Accession	O75881
Other Accession	NM_004820
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	2B11
Calculated MW	58256

Additional Information

Gene ID	9420
Other Names	25-hydroxycholesterol 7-alpha-hydroxylase, Cytochrome P450 7B1, Oxysterol 7-alpha-hydroxylase, CYP7B1
Target/Specificity	CYP7B1 (NP_004811, 203 a.a. ~ 286 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	CYP7B1 Antibody (monoclonal) (M06) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the first reaction in the cholesterol catabolic pathway of extrahepatic tissues, which converts cholesterol to bile acids. This enzyme likely plays a minor role in total bile acid synthesis, but may also be involved in the development of atherosclerosis, neurosteroid metabolism and sex hormone synthesis.


References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. CYP7B1-mediated metabolism of 5 α -androstane-3 α ,17 β -diol (3 α -Adiol): a novel pathway for potential regulation of the cellular levels of androgens and neurosteroids. Pettersson H, et al. Biochim Biophys Acta, 2009 Dec. PMID 19732851. Genetic susceptibility to distinct bladder cancer subphenotypes. Guey LT, et al. Eur Urol, 2010 Feb. PMID 19692168. CYP7B1: one cytochrome P450, two human genetic diseases, and multiple physiological functions. Stiles AR, et al. J Biol Chem, 2009 Oct 16. PMID 19687010.

Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (34.98 KDa) .




Western Blot analysis of CYP7B1 expression in transfected 293T cell line by CYP7B1 monoclonal antibody (M06), clone 2B11.

Lane 1: CYP7B1 transfected lysate (58.256 KDa).
Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to CYP7B1 on formalin-fixed paraffin-embedded human prostate. [antibody concentration 3 μ g/ml]



Detection limit for recombinant GST tagged CYP7B1 is approximately 0.3 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.