

# DPH1 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant DPH1. Catalog # AT1812a

### **Product Information**

Application WB, IF
Primary Accession O9BZG8
Other Accession NM\_001383
Reactivity Human
Host mouse
Clonality monoclonal
Isotype IgG2b Kappa

Clone Names 2C5 Calculated MW 48134

### **Additional Information**

**Gene ID** 1801

Other Names Diphthamide biosynthesis protein 1, DPH1 homolog, HsDph1, Diphthamide

biosynthesis protein 2 homolog-like 1, DPH-like 1, DPH2-like 1, Diphthamide biosynthesis protein 2-like, Ovarian cancer-associated gene 1 protein, DPH1,

DPH2L, DPH2L1, OVCA1

Target/Specificity DPH1 (NP 001374, 216 a.a. ~ 324 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IF~~1:50~200

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** DPH1 Antibody (monoclonal) (M02) is for research use only and not for use in

diagnostic or therapeutic procedures.


## **Background**

Diphthamide is a unique posttranslationally modified histidine found only in translation elongation factor-2 (EEF2; MIM 130610). This modification is conserved from archaebacteria to humans and serves as the target for ADP-ribosylation and inactivation of EEF2 by diphtheria toxin (DT) and Pseudomonas exotoxin A. DPH1 is 1 of several enzymes involved in synthesis of diphthamide in EEF2 (Liu et al., 2004 [PubMed 15485916]).


#### References

Identification of the proteins required for biosynthesis of diphthamide, the target of bacterial ADP-ribosylating toxins on translation elongation factor 2. Liu S, et al. Mol Cell Biol, 2004 Nov. PMID 15485916.Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.Refinement of a 400-kb critical region allows genotypic differentiation between isolated lissencephaly, Miller-Dieker syndrome, and other phenotypes secondary to deletions of 17p13.3. Cardoso C, et al. Am J Hum Genet, 2003 Apr. PMID 12621583.Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932.Cloning, structure, and expression of the mouse Ovca1 gene. Chen CM, et al. Biochem Biophys Res Commun, 2001 Sep 7. PMID 11527402.

### **Images**



DPH1 monoclonal antibody (M02), clone 2C5 Western Blot analysis of DPH1 expression in HeLa ( Cat # L013V1 ).



Immunofluorescence of monoclonal antibody to DPH1 on HeLa cell. [antibody concentration 10 ug/ml]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.