

FES Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant FES.

Catalog # AT2031a

Product Information

| | |
|-------------------|--------------------------|
| Application | WB |
| Primary Accession | P07332 |
| Other Accession | BC035357 |
| Reactivity | Human |
| Host | mouse |
| Clonality | monoclonal |
| Isotype | IgG1 kappa |
| Clone Names | 3A3-1E5 |
| Calculated MW | 93497 |

Additional Information

| | |
|--------------------|---|
| Gene ID | 2242 |
| Other Names | Tyrosine-protein kinase Fes/Fps, Feline sarcoma/Fujinami avian sarcoma oncogene homolog, Proto-oncogene c-Fes, Proto-oncogene c-Fps, p93c-fes, FES, FPS |
| Target/Specificity | FES (AAH35357, 1 a.a. ~ 822 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa. |
| Dilution | WB~~1:500~1000 |
| Format | Clear, colorless solution in phosphate buffered saline, pH 7.2 . |
| Storage | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing. |
| Precautions | FES Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures. |


Background

This gene encodes the human cellular counterpart of a feline sarcoma retrovirus protein with transforming capabilities. The gene product has tyrosine-specific protein kinase activity and that activity is required for maintenance of cellular transformation. Its chromosomal location has linked it to a specific translocation event identified in patients with acute promyelocytic leukemia but it is also involved in normal hematopoiesis as well as growth factor and cytokine receptor signaling. Alternative splicing results in multiple variants encoding different isoforms.


References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.FES kinase participates in KIT-ligand induced chemotaxis. Voisset E, et al. Biochem Biophys Res Commun, 2010 Feb 26. PMID 20117079.Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.Inhibition of endothelial cell chemotaxis toward FGF-2 by gefitinib associates with downregulation of Fes activity. Kanda S, et al. Int J Oncol, 2009 Dec. PMID 19885553.Mutual exclusion of ASXL1 and NPM1 mutations in a series of acute myeloid leukemias. Carbuccia N, et al. Leukemia, 2010 Feb. PMID 19865112.


Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (116.16 kDa) .



FES monoclonal antibody (M01), clone 3A3-1E5 Western Blot analysis of FES expression in HL-60 ((Cat # AT2031a)



Western Blot analysis of FES expression in transfected 293T cell line by FES monoclonal antibody (M01), clone 3A3-1E5.

Lane 1: FES transfected lysate(93 kDa).
Lane 2: Non-transfected lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.