

# FGFR1OP2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant FGFR1OP2. Catalog # AT2038a

#### **Product Information**

Application WB, E
Primary Accession Q9NVK5
Other Accession NM\_015633
Reactivity Human
Host mouse
Clonality monoclonal
Isotype IgG2a Kappa

Clone Names 2G4 Calculated MW 29426

### **Additional Information**

**Gene ID** 26127

Other Names FGFR1 oncogene partner 2, FGFR1OP2

**Target/Specificity** FGFR1OP2 (NP\_056448, 62 a.a. ~ 169 a.a) partial recombinant protein with

GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


**Precautions** FGFR1OP2 Antibody (monoclonal) (M01) is for research use only and not for

use in diagnostic or therapeutic procedures.


#### References

Small cytoskeleton-associated molecule, fibroblast growth factor receptor 1 oncogene partner 2/wound inducible transcript-3.0 (FGFR1OP2/wit3.0), facilitates fibroblast-driven wound closure. Lin A, et al. Am J Pathol, 2010 Jan. PMID 19959814.Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732.A PP2A phosphatase high density interaction network identifies a novel striatin-interacting phosphatase and kinase complex linked to the cerebral cavernous malformation 3 (CCM3) protein. Goudreault M, et al. Mol Cell Proteomics, 2009 Jan. PMID 18782753.Phosphotyrosine profiling identifies the KG-1 cell line as a model for the study of FGFR1 fusions in acute myeloid leukemia. Gu TL, et al. Blood, 2006 Dec 15. PMID 16946300.Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560.

## **Images**




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (37.62 KDa) .



Western Blot analysis of FGFR1OP2 expression in transfected 293T cell line by FGFR1OP2 monoclonal antibody (M01), clone 2G4.

Lane 1: FGFR1OP2 transfected lysate(20.175 KDa). Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged FGFR1OP2 is approximately 0.3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.