

# FUT7 Antibody (monoclonal) (M05)

Mouse monoclonal antibody raised against a partial recombinant FUT7. Catalog # AT2125a

### **Product Information**

**Application** E

Primary Accession
Other Accession
Reactivity
Human
Host
Clonality
Isotype
Q11130
NM\_004479
Human
mouse
Glonality
IgG2a Kappa

Clone Names 1A12 Calculated MW 39239

### **Additional Information**

**Gene ID** 2529

**Other Names** Alpha-(1, 3)-fucosyltransferase 7, 241-, Fucosyltransferase 7,

Fucosyltransferase VII, Fuc-TVII, FucT-VII, Galactoside 3-L-fucosyltransferase,

Selectin ligand synthase, FUT7

**Target/Specificity** FUT7 (NP\_004470, 262 a.a. ~ 342 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

**Dilution** E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** FUT7 Antibody (monoclonal) (M05) is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Background**


The protein encoded by this gene is a Golgi stack membrane protein that is involved in the creation of sialyl-Lewis X antigens. The encoded protein can direct the synthesis of the E-selectin-binding sialyl-Lewis X moiety.

#### References

Association of genetic variants with hemorrhagic stroke in Japanese individuals. Yoshida T, et al. Int J Mol Med, 2010 Apr. PMID 20198315. Assessment of a polymorphism of SDK1 with hypertension in Japanese

Individuals. Oguri M, et al. Am J Hypertens, 2010 Jan. PMID 19851296.alpha1,3 fucosyltransferase-VII up-regulates the mRNA of alpha5 integrin and its biological function. Wang QY, et al. J Cell Biochem, 2008 Aug 15. PMID 18452157.Overexpression of fucosyltransferase VII (FUT7) promotes embryo adhesion and implantation. Zhang Y, et al. Fertil Steril, 2009 Mar. PMID 18402946.Evaluation of urinary CA19-9 levels in bladder cancer patients classified according to the combinations of Lewis and Secretor blood group genotypes. Nagao K, et al. Int J Urol, 2007 Sep. PMID 17760744.

### **Images**



Detection limit for recombinant GST tagged FUT7 is approximately 0.1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.