

FXR1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant FXR1. Catalog # AT2127a

Product Information

Application WB, IHC, E **Primary Accession** P51114 **Other Accession** BC028983 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 kappa **Clone Names** 2G11 **Calculated MW** 69721

Additional Information

Gene ID 8087

Other Names Fragile X mental retardation syndrome-related protein 1, hFXR1p, FXR1

Target/Specificity FXR1 (AAH28983, 121 a.a. ~ 220 a.a) partial recombinant protein with GST tag.

MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 IHC~~1:100~500 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions FXR1 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.

Background


The protein encoded by this gene is an RNA binding protein that interacts with the functionally-similar proteins FMR1 and FXR2. These proteins shuttle between the nucleus and cytoplasm and associate with polyribosomes, predominantly with the 60S ribosomal subunit. Three transcript variants encoding different isoforms have been found for this gene.

References


Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. International Schizophrenia Consortium, et al. Nature, 2009 Aug 6. PMID 19571811. Discrimination of common and unique RNA-binding activities among Fragile X mental retardation protein paralogs. Darnell JC, et al. Hum Mol

Genet, 2009 Sep 1. PMID 19487368.Alteration of expression of muscle specific isoforms of the fragile X related protein 1 (FXR1P) in facioscapulohumeral muscular dystrophy patients. Davidovic L, et al. J Med Genet, 2008 Oct. PMID 18628314.Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.AU-rich-element-mediated upregulation of translation by FXR1 and Argonaute 2. Vasudevan S, et al. Cell, 2007 Mar 23. PMID 17382880.

Images




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.63 KDa) .




Western Blot analysis of FXR1 expression in transfected 293T cell line by FXR1 monoclonal antibody (M01), clone 2G11.

Lane 1: FXR1 transfected lysate(61 KDa). Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to FXR1 on formalin-fixed paraffin-embedded human colon adenocarcinoma. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged FXR1 is approximately 0.3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.