

# HIPK1 Antibody (monoclonal) (M07)

Mouse monoclonal antibody raised against a partial recombinant HIPK1. Catalog # AT2373a

#### **Product Information**

ApplicationWB, IF, EPrimary AccessionQ86Z02Other AccessionBC028408ReactivityHumanHostmouseClonalitymonoclonalIsotypeIgG2a Kappa

Clone Names 40 Calculated MW 130843

### **Additional Information**

**Gene ID** 204851

**Other Names** Homeodomain-interacting protein kinase 1, Nuclear body-associated kinase 2,

HIPK1, KIAA0630, MYAK, NBAK2

Target/Specificity HIPK1 (AAH28408, 330 a.a. ~ 430 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IF~~1:50~200 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** HIPK1 Antibody (monoclonal) (M07) is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Background**


The protein encoded by this gene belongs to the Ser/Thr family of protein kinases and HIPK subfamily. It phosphorylates homeodomain transcription factors and may also function as a co-repressor for homeodomain transcription factors. Alternative splicing results in four transcript variants encoding four distinct isoforms.

### References


HIPK1 interacts with c-Myb and modulates its activity through phosphorylation. Matre V, et al. Biochem Biophys Res Commun, 2009 Oct 9. PMID 19646965.Toward a confocal subcellular atlas of the human

proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.Roles of HIPK1 and HIPK2 in AML1-and p300-dependent transcription, hematopoiesis and blood vessel formation. Aikawa Y, et al. EMBO J, 2006 Sep 6. PMID 16917507.Tumor necrosis factor alpha-induced desumoylation and cytoplasmic translocation of homeodomain-interacting protein kinase 1 are critical for apoptosis signal-regulating kinase 1-JNK/p38 activation. Li X, et al. J Biol Chem, 2005 Apr 15. PMID 15701637.The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.


### **Images**



HIPK1 monoclonal antibody (M07), clone 4E1 Western Blot analysis of HIPK1 expression in HeLa ( Cat # L013V1 ).



Immunofluorescence of monoclonal antibody to HIPK1 on HeLa cell. [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged HIPK1 is approximately 0.3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.