

HMGA2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant HMGA2.

Catalog # AT2383a

Product Information

Application	WB, IF, E
Primary Accession	P52926
Other Accession	NM_003483
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 Kappa
Clone Names	2D10
Calculated MW	11832

Additional Information

Gene ID	8091
Other Names	High mobility group protein HMGI-C, High mobility group AT-hook protein 2, HMGA2, HMGIC
Target/Specificity	HMGA2 (NP_003474, 1 a.a. ~ 92 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IF~~1:50~200 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	HMGA2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

This gene encodes a protein that belongs to the non-histone chromosomal high mobility group (HMG) protein family. HMG proteins function as architectural factors and are essential components of the enhancosome. This protein contains structural DNA-binding domains and may act as a transcriptional regulating factor. Identification of the deletion, amplification, and rearrangement of this gene that are associated with myxoid liposarcoma suggests a role in adipogenesis and mesenchymal differentiation. A gene knock out study of the mouse counterpart demonstrated that this gene is involved in diet-induced obesity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.


References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Gamma-Radiation Sensitivity and Polymorphisms in RAD51L1 Modulate Glioma Risk. Liu Y, et al. Carcinogenesis, 2010 Jul 7. PMID 20610542. HMGA2 is a sensitive but not specific immunohistochemical marker of vulvovaginal aggressive angiomyxoma. McCluggage WG, et al. Am J Surg Pathol, 2010 Jul. PMID 20551826. HMGA2 and the p19Arf-TP53-CDKN1A axis: a delicate balance in the growth of uterine leiomyomas. Markowski DN, et al. Genes Chromosomes Cancer, 2010 Aug. PMID 20544840. HMGA1 and HMGA2 rearrangements in mass-forming endometriosis. Medeiros F, et al. Genes Chromosomes Cancer, 2010 Jul. PMID 20461755.


Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.86 kDa) .



Immunofluorescence of monoclonal antibody to HMGA2 on HeLa cell . [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged HMGA2 is approximately 0.1 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.