

ICOSLG Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant ICOSLG.

Catalog # AT2470a

Product Information


Application	WB, E
Primary Accession	O75144
Other Accession	BC064637
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 kappa
Clone Names	2D12
Calculated MW	33349

Additional Information


Gene ID	23308
Other Names	ICOS ligand, B7 homolog 2, B7-H2, B7-like protein GI50, B7-related protein 1, B7RP-1, CD275, ICOSLG, B7H2, B7RP1, ICOSL, KIAA0653
Target/Specificity	ICOSLG (AAH64637, 20 a.a. ~ 302 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	ICOSLG Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

References


An approach based on a genome-wide association study reveals candidate loci for narcolepsy. Shimada M, et al. Hum Genet, 2010 Oct. PMID 20677014. Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Impact of female cigarette smoking on circulating B cells in vivo: the suppressed ICOSLG, TCF3, and VCAM1 gene functional network may inhibit normal cell function. Pan F, et al. Immunogenetics, 2010 Apr. PMID 20217071. Multiple common variants for celiac disease influencing immune gene expression. Dubois PC, et al. Nat Genet, 2010 Apr. PMID 20190752. Comparative genetic analysis of inflammatory bowel disease and type 1 diabetes implicates multiple loci with opposite effects. Wang K, et al. Hum Mol Genet, 2010 May 15. PMID 20176734.



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (56.87 KDa) .



ICOSLG monoclonal antibody (M01), clone 2D12 Western Blot analysis of ICOSLG expression in HL-60 ((Cat # AT2470a)



Detection limit for recombinant GST tagged ICOSLG is approximately 0.03ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.