

ISG20 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant ISG20. Catalog # AT2557a

Product Information

Application WB, IF, E **Primary Accession Q96AZ6** Other Accession BC007922 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 kappa **Clone Names** 1B2-3C9 Calculated MW 20363

Additional Information

Gene ID 3669

Other Names Interferon-stimulated gene 20 kDa protein, Estrogen-regulated transcript 45

protein, Promyelocytic leukemia nuclear body-associated protein ISG20,

ISG20, HEM45

Target/Specificity ISG20 (AAH07922.1, 1 a.a. ~ 181 a.a) full-length recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 IF~~1:50~200 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


Precautions ISG20 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.


References

1.Antiviral activities of ISG20 in positive-strand RNA virus infections.Zhou Z, Wang N, Woodson SE, Dong Q, Wang J, Liang Y, Rijnbrand R, Wei L, Nichols JE, Guo JT, Holbrook MR, Lemon SM, Li K.Virology. 2010 Oct 29. [Epub ahead of print]2.Intracellular activation of interferon regulatory factor-1 by nanobodies to the multi-functional (Mf1) domain.Moeller A, Pion E, Narayan V, Ball KL.J Biol Chem. 2010 Sep 3. [Epub ahead of print]

Images




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (45.65 KDa).




Western Blot analysis of ISG20 expression in transfected 293T cell line by ISG20 monoclonal antibody (M01), clone 1B2-3C9.

Lane 1: ISG20 transfected lysate(20.4 KDa). Lane 2: Non-transfected lysate.



Immunofluorescence of monoclonal antibody to ISG20 on HeLa cell . [antibody concentration 10 ug/ml]



Recombinant ProteinConcentration(ng/ml)

Detection limit for recombinant GST tagged ISG20 is approximately 0.1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.