

## MAZ Antibody (monoclonal) (M05)

Mouse monoclonal antibody raised against a partial recombinant MAZ.

Catalog # AT2809a

### Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P56270</a>
<b>Other Accession</b>	<a href="#">NM_002383</a>
<b>Reactivity</b>	Human
<b>Host</b>	mouse
<b>Clonality</b>	monoclonal
<b>Isotype</b>	IgG2a Kappa
<b>Clone Names</b>	6G2
<b>Calculated MW</b>	48608

### Additional Information

---


<b>Gene ID</b>	4150
<b>Other Names</b>	Myc-associated zinc finger protein, MAZI, Pur-1, Purine-binding transcription factor, Serum amyloid A-activating factor-1, SAF-1, Transcription factor Zif87, ZF87, Zinc finger protein 801, MAZ, ZNF801
<b>Target/Specificity</b>	MAZ (NP_002374, 332 a.a. ~ 440 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
<b>Dilution</b>	WB~~1:500~1000 E~~N/A
<b>Format</b>	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
<b>Storage</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Precautions</b>	MAZ Antibody (monoclonal) (M05) is for research use only and not for use in diagnostic or therapeutic procedures.

### References


---

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732.SAF-3, a novel splice variant of the SAF-1/MAZ/Pur-1 family, is expressed during inflammation. Ray A, et al. FEBS J, 2009 Aug. PMID 19583771.Association and mutation analyses of 16p11.2 autism candidate genes. Kumar RA, et al. PLoS One, 2009. PMID 19242545.Gastrin activates paracrine networks leading to induction of PAI-2 via MAZ and ASC-1. Almeida-Vega S, et al. Am J Physiol Gastrointest Liver Physiol, 2009 Feb. PMID 19074642.Transcriptional synergy mediated by SAF-1 and AP-1: critical role of N-terminal polyalanine and two zinc finger domains of SAF-1. Kumar D, et al. J Biol Chem, 2009 Jan 16. PMID 19028685.

## Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 KDa) .



Detection limit for recombinant GST tagged MAZ is approximately 0.1 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.