

# MEIS2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant MEIS2. Catalog # AT2841a

### **Product Information**

**Application** WB, IHC, IF, E **Primary Accession** <u>014770</u> **Other Accession** BC001516 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 kappa **Clone Names** 1H4 **Calculated MW** 51790

#### **Additional Information**

**Gene ID** 4212

Other Names Homeobox protein Meis2, Meis1-related protein 1, MEIS2, MRG1

Target/Specificity MEIS2 (AAH01516, 1 a.a. ~ 381 a.a) full-length recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** MEIS2 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.

# **Background**


This gene encodes a homeobox protein belonging to the TALE ('three amino acid loop extension') family of homeodomain-containing proteins. TALE homeobox proteins are highly conserved transcription regulators, and several members have been shown to be essential contributors to developmental programs. Multiple transcript variants encoding distinct isoforms have been described for this gene.

### References


1.Cooperative transcriptional activation by klf4, meis2, and pbx1.Bjerke GA, Hyman-Walsh C, Wotton D.Mol Cell Biol. 2011 Sep;31(18):3723-33. Epub 2011 Jul 11.2.A neuronal migratory pathway crossing from diencephalon to telencephalon populates amygdala nuclei.Garcia-Moreno F, Pedraza M, Di Giovannantonio

LG, Di Salvio M, Lopez-Mascaraque L, Simeone A, De Carlos JA.Nat Neurosci. 2010 Jun;13(6):680-9. Epub 2010 May 23.


## **Images**




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (67.65 KDa).



MEIS2 monoclonal antibody (M01), clone 1H4 Western Blot analysis of MEIS2 expression in HeLa ( (Cat # AT2841a )



MEIS2 monoclonal antibody (M01), clone 1H4. Western Blot analysis of MEIS2 expression in K-562 ( (Cat # AT2841a )




Western Blot analysis of MEIS2 expression in transfected 293T cell line by MEIS2 monoclonal antibody (M01), clone 1H4.


Lane 1: MEIS2 transfected lysate(51.1 KDa).

Lane 2: Non-transfected lysate.


Immunoperoxidase of monoclonal antibody to MEIS2 on formalin-fixed paraffin-embedded human spleen tissue.



[antibody concentration 1 ug/ml]



Immunofluorescence of monoclonal antibody to MEIS2 on HeLa cell. [antibody concentration 10 ug/ml]



Recombinant ProteinConcentration(ng/ml)

Detection limit for recombinant GST tagged MEIS2 is approximately 0.03ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.