

# MGC10540 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant MGC10540. Catalog # AT2857a

### **Product Information**

**Application** WB, IF, E **Primary Accession** Q9BRG1 **Other Accession** BC006282 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 kappa **Clone Names** 2E5-2B9 Calculated MW 20748

#### **Additional Information**

**Gene ID** 84313

**Other Names** Vacuolar protein-sorting-associated protein 25, hVps25, Dermal

papilla-derived protein 9, ELL-associated protein of 20 kDa, ESCRT-II complex

subunit VPS25, VPS25, DERP9, EAP20

Target/Specificity MGC10540 (AAH06282, 1 a.a. ~ 176 a.a) full-length recombinant protein with

GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IF~~1:50~200 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** MGC10540 Antibody (monoclonal) (M01) is for research use only and not for

use in diagnostic or therapeutic procedures.

## **Background**


VPS25, VPS36 (MIM 610903), and SNF8 (MIM 610904) form ESCRT-II (endosomal sorting complex required for transport II), a complex involved in endocytosis of ubiquitinated membrane proteins. VPS25, VPS36, and SNF8 are also associated in a multiprotein complex with RNA polymerase II elongation factor (ELL; MIM 600284) (Slagsvold et al., 2005 [PubMed 15755741]; Kamura et al., 2001 [PubMed 11278625]).

### References


Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes

REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.Structure and function of the ESCRT-II-III interface in multivesicular body biogenesis. Im YJ, et al. Dev Cell, 2009 Aug. PMID 19686684.Degradation of endocytosed epidermal growth factor and virally ubiquitinated major histocompatibility complex class I is independent of mammalian ESCRTII. Bowers K, et al. J Biol Chem, 2006 Feb 24. PMID 16371348.Towards a proteome-scale map of the human protein-protein interaction network. Rual JF, et al. Nature, 2005 Oct 20. PMID 16189514.


### **Images**




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (45.1 KDa).



MGC10540 monoclonal antibody (M01), clone 2E5-2B9 Western Blot analysis of MGC10540 expression in Hela S3 NE ( (Cat # AT2857a )



Immunofluorescence of monoclonal antibody to VPS25 on HeLa cell. [antibody concentration 10 ug/ml]



Recombinant ProteinConcentration(ng/ml)

Detection limit for recombinant GST tagged VPS25 is approximately 0.03ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.