10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MSLN Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant MSLN.
Catalog # AT2917a

Product Information

Application	WB, IP, E
Primary Accession	Q13421
Other Accession	NM_005823
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG2b Kappa
Clone Names	1A10
Calculated MW	68070

Additional Information

Gene ID	10232
Other Names	Mesothelin, CAK1 antigen, Pre-pro-megakaryocyte-potentiating factor, Megakaryocyte-potentiating factor, MPF, Mesothelin, cleaved form, MSLN, MPF
Target/Specificity	MSLN (NP_005814, 464 a.a. ~ 563 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IP~~N/A E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	MSLN Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

This gene encodes a precursor protein that is cleaved into two products, megakaryocyte potentiating factor and mesothelin. Megakaryocyte potentiation factor functions as a cytokine that can stimulate colony formation in bone marrow megakaryocytes. Mesothelian is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed in epithelial mesotheliomas, ovarian cancers and in specific squamous cell carcinomas. Alternative splicing results in multiple transcript variants.


References

Mesothelin (MSLN) promoter is hypomethylated in malignant mesothelioma, but its expression is not associated with methylation status of the promoter. Tan K, et al. Hum Pathol, 2010 Sep. PMID 20573372.A Large-scale genetic association study of esophageal adenocarcinoma risk. Liu CY, et al. Carcinogenesis, 2010 Jul. PMID 20453000.Mesothelin gene expression and promoter methylation/hypomethylation in gynecological tumors. Obulhasim G, et al. Eur J Gynaecol Oncol, 2010. PMID 20349783.Soluble mesothelin related peptides (SMRP) and osteopontin as protein biomarkers for malignant mesothelioma: analytical validation of ELISA based assays and characterization at mRNA and protein levels. Rai AJ, et al. Clin Chem Lab Med, 2010 Feb. PMID 20131968.Polymorphisms in the putative micro-RNA-binding sites of mesothelin gene are associated with serum levels of mesothelin-related protein. Cristaudo A, et al. Occup Environ Med, 2010 Apr. PMID 19858537.


Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.74 KDa) .



Immunoprecipitation of MSLN transfected lysate using anti-MSLN monoclonal antibody and Protein A Magnetic Bead (U0007), and immunoblotted with MSLN MaxPab rabbit polyclonal antibody.



Detection limit for recombinant GST tagged MSLN is approximately 1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.