10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MT Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant MT.
Catalog # AT2921a

Product Information

Application	WB, IHC, E
Primary Accession	Q8IVS2
Other Accession	NM_173467
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	2F2
Calculated MW	42962

Additional Information

Gene ID	27349
Other Names	Malonyl-CoA-acyl carrier protein transacylase, mitochondrial, MCT, Mitochondrial malonyl CoA:ACP acyltransferase, Mitochondrial malonyltransferase, [Acyl-carrier-protein] malonyltransferase, MCAT, MT
Target/Specificity	MT (NP_775738, 291 a.a. ~ 390 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	MT Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

The protein encoded by this gene is found exclusively in the mitochondrion, where it catalyzes the transfer of a malonyl group from malonyl-CoA to the mitochondrial acyl carrier protein. The encoded protein may be part of a fatty acid synthase complex that is more like the type II prokaryotic and plastid complexes rather than the type I human cytosolic complex. Two transcript variants encoding different isoforms have been found for this gene.


References

Prostate cancer risk-associated variants reported from genome-wide association studies: Meta-analysis and their contribution to genetic Variation. Kim ST, et al. Prostate, 2010 Jun 16. PMID 20564319.Identification of seven new prostate cancer susceptibility loci through a genome-wide association study. Eeles RA, et al. Nat Genet, 2009 Oct. PMID 19767753.Identifying leukocyte gene expression patterns associated with plasma lipid levels in human subjects. Ma J, et al. Atherosclerosis, 2007 Mar. PMID 16806233.Exercise training decreases the concentration of malonyl-CoA and increases the expression and activity of malonyl-CoA decarboxylase in human muscle. Kuhl JE, et al. Am J Physiol Endocrinol Metab, 2006 Jun. PMID 16434556.Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560.


Images




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.74 KDa) .



MT monoclonal antibody (M01), clone 2F2 Western Blot analysis of MT expression in HeLa ((Cat # AT2921a)



Immunoperoxidase of monoclonal antibody to MT on formalin-fixed paraffin-embedded human colon. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged MT is approximately 1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.