

MYL1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant MYL1.

Catalog # AT2951a

Product Information

Application	WB, IHC, E
Primary Accession	P05976
Other Accession	NM_079422
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	2D9
Calculated MW	21145

Additional Information

Gene ID	4632
Other Names	Myosin light chain 1/3, skeletal muscle isoform, MLC1/MLC3, MLC1F/MLC3F, Myosin light chain alkali 1/2, Myosin light chain A1/A2, MYL1
Target/Specificity	MYL1 (NP_524146, 1 a.a. ~ 80 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	MYL1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

Background


Myosin is a hexameric ATPase cellular motor protein. It is composed of two heavy chains, two nonphosphorylatable alkali light chains, and two phosphorylatable regulatory light chains. This gene encodes a myosin alkali light chain expressed in fast skeletal muscle. Two transcript variants have been identified for this gene.

References


Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560. The status,

quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932. Interaction of ADP-ribosylated actin with actin binding proteins. Ballweber E, et al. FEBS Lett, 2001 Nov 9. PMID 11707283. A nonerythroid isoform of protein 4.1R interacts with components of the contractile apparatus in skeletal myofibers. Kontogianni-Konstantopoulos A, et al. Mol Biol Cell, 2000 Nov. PMID 11071908.

Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (34.54 KDa) .




Western Blot analysis of MYL1 expression in transfected 293T cell line by MYL1 monoclonal antibody (M01), clone 2D9.

Lane 1: MYL1 transfected lysate(21.1 KDa).
Lane 2: Non-transfected lysate.




Immunoperoxidase of monoclonal antibody to MYL1 on formalin-fixed paraffin-embedded human spleen. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged MYL1 is approximately 3ng/ml as a capture antibody.

Western blot analysis of MYL1 over-expressed 293 cell line, cotransfected with MYL1 Validated Chimera RNAi (



(Cat # AT2951a)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.