10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MYOG Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant MYOG.
Catalog # AT2963a

Product Information

Application	WB, E
Primary Accession	P15173
Other Accession	BC053899
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 kappa
Clone Names	2B7
Calculated MW	25037

Additional Information

Gene ID	4656
Other Names	Myogenin, Class C basic helix-loop-helix protein 3, bHLHc3, Myogenic factor 4, Myf-4, MYOG, BHLHC3, MYF4
Target/Specificity	MYOG (AAH53899, 1 a.a. ~ 224 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	MYOG Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

Background


Myogenin is a muscle-specific transcription factor that can induce myogenesis in a variety of cell types in tissue culture. It is a member of a large family of proteins related by sequence homology, the helix-loop-helix (HLH) proteins. It is essential for the development of functional skeletal muscle.

References


CARM1 activates myogenin gene via PCAF in the early differentiation of TPA-induced rhabdomyosarcoma-derived cells. Gao X, et al. J Cell Biochem, 2010 May. PMID 20213728.Decreased Jun-D and myogenin expression in muscle wasting of human cachexia. Ramamoorthy S, et al. Am J Physiol

Endocrinol Metab, 2009 Aug. PMID 19470832.High-density association study of 383 candidate genes for volumetric BMD at the femoral neck and lumbar spine among older men. Yerges LM, et al. J Bone Miner Res, 2009 Dec. PMID 19453261.Opposing control of rhabdomyosarcoma growth and differentiation by myogenin and interleukin 4. Nanni P, et al. Mol Cancer Ther, 2009 Apr. PMID 19372547.SMD and NMD are competitive pathways that contribute to myogenesis: effects on PAX3 and myogenin mRNAs. Gong C, et al. Genes Dev, 2009 Jan 1. PMID 19095803.

Images




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (50.38 KDa) .




Western Blot analysis of MYOG expression in transfected 293T cell line by MYOG monoclonal antibody (M01), clone 2B7.

Lane 1: MYOG transfected lysate(25 KDa).
Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged MYOG is approximately 0.03ng/ml as a capture antibody.



Western blot analysis of MYOG over-expressed 293 cell line, cotransfected with MYOG Validated Chimera RNAi ((Cat # AT2963a)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.