

NDUFS4 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant NDUFS4.

Catalog # AT3007a

Product Information

Application	WB, IHC, E
Primary Accession	O43181
Other Accession	NM_002495
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	1A1
Calculated MW	20108

Additional Information

Gene ID	4724
Other Names	NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial, Complex I-18 kDa, CI-18 kDa, Complex I-AQDQ, CI-AQDQ, NADH-ubiquinone oxidoreductase 18 kDa subunit, NDUFS4
Target/Specificity	NDUFS4 (NP_002486, 66 a.a. ~ 175 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	NDUFS4 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), or NADH:ubiquinone oxidoreductase, the first multi-subunit enzyme complex of the mitochondrial respiratory chain. Complex I plays a vital role in cellular ATP production, the primary source of energy for many crucial processes in living cells. It removes electrons from NADH and passes them by a series of different protein-coupled redox centers to the electron acceptor ubiquinone. In well-coupled mitochondria, the electron flux leads to ATP generation via the building of a proton gradient across the inner membrane. Complex I is composed of at least 41 subunits, of which 7 are encoded by the mitochondrial genome and the remainder by nuclear genes.


References

NDUFS4 mutations cause Leigh syndrome with predominant brainstem involvement. Leshinsky-Silver E, et al. Mol Genet Metab, 2009 Jul. PMID 19364667. Association study between single-nucleotide polymorphisms in 199 drug-related genes and commonly measured quantitative traits of 752 healthy Japanese subjects. Saito A, et al. J Hum Genet, 2009 Jun. PMID 19343046. Polymorphisms in mitochondrial genes and prostate cancer risk. Wang L, et al. Cancer Epidemiol Biomarkers Prev, 2008 Dec. PMID 19064571. Oxidative stress, telomere length and biomarkers of physical aging in a cohort aged 79 years from the 1932 Scottish Mental Survey. Starr JM, et al. Mech Ageing Dev, 2008 Dec. PMID 18977241. The regulation of PTC containing transcripts of the human NDUFS4 gene of complex I of respiratory chain and the impact of pathological mutations. Panelli D, et al. Biochimie, 2008 Oct. PMID 18555024.


Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .





NDUFS4 monoclonal antibody (M01), clone 1A1 Western Blot analysis of NDUFS4 expression in A-431 ((Cat # AT3007a)




Western Blot analysis of NDUFS4 expression in transfected 293T cell line by NDUFS4 monoclonal antibody (M01), clone 1A1.

Lane 1: NDUFS4 transfected lysate(20.1 KDa).
Lane 2: Non-transfected lysate.

Immunoperoxidase of monoclonal antibody to NDUFS4 on formalin-fixed paraffin-embedded human colon. [antibody concentration 1 ug/ml]



Detection limit for recombinant GST tagged NDUF54 is approximately 0.3ng/ml as a capture antibody.



Western blot analysis of NDUF54 over-expressed 293 cell line, cotransfected with NDUF54 Validated Chimera RNAi (Cat # AT3007a)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.