

NEFH Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant NEFH.

Catalog # AT3010a

Product Information

Application	WB, E
Primary Accession	P12036
Other Accession	NM_021076
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	2E9
Calculated MW	111838

Additional Information

Gene ID	4744
Other Names	Neurofilament heavy polypeptide, NF-H, 200 kDa neurofilament protein, Neurofilament triplet H protein, NEFH, KIAA0845, NFH
Target/Specificity	NEFH (NP_066554, 263 a.a. ~ 363 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	NEFH Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

Background


Neurofilaments are type IV intermediate filament heteropolymers composed of light, medium, and heavy chains. Neurofilaments comprise the axoskeleton and functionally maintain neuronal caliber. They may also play a role in intracellular transport to axons and dendrites. This gene encodes the heavy neurofilament protein. This protein is commonly used as a biomarker of neuronal damage and susceptibility to amyotrophic lateral sclerosis (ALS) has been associated with mutations in this gene.

References


A genome-wide association study of carotid atherosclerosis in HIV-infected men. Shrestha S, et al. AIDS,

2010 Feb 20. PMID 20009918. Comparative study of CSF neurofilaments in HTLV-1-associated myelopathy/tropical spastic paraparesis and other neurological disorders. Alberti C, et al. AIDS Res Hum Retroviruses, 2009 Aug. PMID 19678766. Phosphorylated neurofilament heavy chain is a marker of neurodegeneration in Leber hereditary optic neuropathy (LHON). Guy J, et al. Mol Vis, 2008. PMID 19104679. Polymorphisms in genes involved in neurodevelopment may be associated with altered brain morphology in schizophrenia: preliminary evidence. Gregorio SP, et al. Psychiatry Res, 2009 Jan 30. PMID 19054571. Pin1-dependent prolyl isomerization modulates the stress-induced phosphorylation of high molecular weight neurofilament protein. Rudrabhatla P, et al. J Biol Chem, 2008 Sep 26. PMID 18635547.


Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.85 KDa) .



NEFH monoclonal antibody (M01), clone 2E8 Western Blot analysis of NEFH expression in HeLa (Cat # AT3010a)



Detection limit for recombinant GST tagged NEFH is approximately 0.03ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.