

## PARC Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant PARC.

Catalog # AT3180a

### Product Information

---

|                          |                           |
|--------------------------|---------------------------|
| <b>Application</b>       | WB                        |
| <b>Primary Accession</b> | <a href="#">Q8IWT3</a>    |
| <b>Other Accession</b>   | <a href="#">NM_015089</a> |
| <b>Reactivity</b>        | Human                     |
| <b>Host</b>              | mouse                     |
| <b>Clonality</b>         | monoclonal                |
| <b>Isotype</b>           | IgG2a Kappa               |
| <b>Clone Names</b>       | 3F7                       |
| <b>Calculated MW</b>     | 281229                    |

### Additional Information

---

|                           |                                                                                                                          |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------|
| <b>Gene ID</b>            | 23113                                                                                                                    |
| <b>Other Names</b>        | Cullin-9, CUL-9, Ubch7-associated protein 1, p53-associated parkin-like cytoplasmic protein, CUL9, H7AP1, KIAA0708, PARC |
| <b>Target/Specificity</b> | PARC (NP_055904, 918 a.a. ~ 1025 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.       |
| <b>Dilution</b>           | WB~~1:500~1000                                                                                                           |
| <b>Format</b>             | Clear, colorless solution in phosphate buffered saline, pH 7.2 .                                                         |
| <b>Storage</b>            | Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.                                                 |
| <b>Precautions</b>        | PARC Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.       |


### References

---

Association of genetic variants with hemorrhagic stroke in Japanese individuals. Yoshida T, et al. Int J Mol Med, 2010 Apr. PMID 20198315. Assessment of a polymorphism of SDK1 with hypertension in Japanese Individuals. Oguri M, et al. Am J Hypertens, 2010 Jan. PMID 19851296. Activation of p53-dependent responses in tumor cells treated with a PARC-interacting peptide. Vitali R, et al. Biochem Biophys Res Commun, 2008 Apr 4. PMID 18230339. Systematic identification of SH3 domain-mediated human protein-protein interactions by peptide array target screening. Wu C, et al. Proteomics, 2007 Jun. PMID 17474147. Hyperubiquitylation of wild-type p53 contributes to cytoplasmic sequestration in neuroblastoma. Becker K, et al. Cell Death Differ, 2007 Jul. PMID 17380154.

## Images

---



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.62 KDa) .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.