

PARK2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant PARK2. Catalog # AT3182a

Product Information

Application WB, IF, E **Primary Accession** <u>060260</u> **Other Accession** BC022014 Reactivity Human Host mouse Clonality monoclonal Isotype IgG3 Kappa **Clone Names** 1H4 Calculated MW 51641

Additional Information

Gene ID 5071

Other Names E3 ubiquitin-protein ligase parkin, 632-, Parkinson juvenile disease protein 2,

Parkinson disease protein 2, PARK2, PRKN

Target/Specificity PARK2 (AAH22014, 288 a.a. ~ 387 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 IF~~1:50~200 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions PARK2 Antibody (monoclonal) (M01) is for research use only and not for use

in diagnostic or therapeutic procedures.


Background

The precise function of this gene is unknown; however, the encoded protein is a component of a multiprotein E3 ubiquitin ligase complex that mediates the targeting of substrate proteins for proteasomal degradation. Mutations in this gene are known to cause Parkinson disease and autosomal recessive juvenile Parkinson disease. Alternative splicing of this gene produces multiple transcript variants encoding distinct isoforms. Additional splice variants of this gene have been described but currently lack transcript support.


References

^{1.}Regional and cellular localisation of Parkin Co-Regulated Gene in developing and adult mouse brain. Brody


Images




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.74 KDa) .



PARK2 monoclonal antibody (M01), clone 1H4 Western Blot analysis of PARK2 expression in Jurkat ((Cat # AT3182a)



Immunofluorescence of monoclonal antibody to PARK2 on HeLa cell. [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged PARK2 is 0.1 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.