

# PBX1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant PBX1. Catalog # AT3200a

#### **Product Information**

Application WB, IHC, IF, E
Primary Accession P40424
Other Accession NM\_002585
Reactivity Human
Host mouse
Clonality monoclonal
Isotype IgG2a Kappa

Clone Names 4A2 Calculated MW 46626

### **Additional Information**

**Gene ID** 5087

Other Names Pre-B-cell leukemia transcription factor 1, Homeobox protein PBX1,

Homeobox protein PRL, PBX1, PRL

**Target/Specificity** PBX1 (NP\_002576, 213 a.a. ~ 321 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


**Precautions** PBX1 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.


### References

1.PBX1 Genomic Pioneer Function Drives ER alpha Signaling Underlying Progression in Breast Cancer.Magnani L, Ballantyne EB, Zhang X, Lupien M.PLoS Genet. 2011 Nov;7(11):e1002368. Epub 2011 Nov 17.2.Cooperative transcriptional activation by klf4, meis2, and pbx1.Bjerke GA, Hyman-Walsh C, Wotton D.Mol Cell Biol. 2011 Sep;31(18):3723-33. Epub 2011 Jul 11.3.Pre-B-cell leukemia homeobox 1 (PBX1) shows functional and possible genetic association with bone mineral density variation.Cheung CL, Chan BY, Chan V, Ikegawa S, Kou I, Ngai H, Smith D, Luk KD, Huang QY, Mori S, Sham PC, Kung AW.Hum Mol Genet. 2009 Feb 15;18(4):679-87. Epub 2008 Dec 8.


## **Images**




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (37.73 KDa).




PBX1 monoclonal antibody (M01), clone 4A2 Western Blot analysis of PBX1 expression in Hela S3 NE ( (Cat # AT3200a )



Immunoperoxidase of monoclonal antibody to PBX1 on formalin-fixed paraffin-embedded human salivary gland. [antibody concentration 3 ug/ml]



Immunofluorescence of monoclonal antibody to PBX1 on HeLa cell. [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged PBX1 is approximately 0.03ng/ml as a capture antibody.