

PCDH7 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant PCDH7.

Catalog # AT3215a

Product Information

Application	WB, E
Primary Accession	O60245
Other Accession	NM_002589
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	2D7
Calculated MW	116071

Additional Information

Gene ID	5099
Other Names	Protocadherin-7, Brain-heart protocadherin, BH-Pcdh, PCDH7, BHPCDH
Target/Specificity	PCDH7 (NP_002580, 31 a.a. ~ 124 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	PCDH7 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

Background


This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily. The gene encodes a protein with an extracellular domain containing 7 cadherin repeats. The gene product is an integral membrane protein that is thought to function in cell-cell recognition and adhesion. Alternative splicing yields isoforms with unique cytoplasmic tails.

References


Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.Genome-wide analysis of survival in early-stage non-small-cell lung cancer. Huang YT, et al. J Clin Oncol, 2009 Jun 1. PMID 19414679.Global, in vivo, and

site-specific phosphorylation dynamics in signaling networks. Olsen JV, et al. Cell, 2006 Nov 3. PMID 17081983. Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. Rush J, et al. Nat Biotechnol, 2005 Jan. PMID 15592455. Large-scale characterization of HeLa cell nuclear phosphoproteins. Beausoleil SA, et al. Proc Natl Acad Sci U S A, 2004 Aug 17. PMID 15302935.

Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.08 KDa) .



Detection limit for recombinant GST tagged PCDH7 is approximately 0.3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.