

PDE1B Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant PDE1B. Catalog # AT3251a

Product Information

ApplicationWB, EPrimary AccessionQ01064Other AccessionBC032226ReactivityHumanHostmouseClonalitymonoclonalIsotypeIgG2a Kappa

Clone Names 5B6 Calculated MW 61380

Additional Information

Gene ID 5153

Other Names Calcium/calmodulin-dependent 3', 5'-cyclic nucleotide phosphodiesterase 1B,

Cam-PDE 1B, 63 kDa Cam-PDE, PDE1B, PDE1B1, PDES1B

Target/Specificity PDE1B (AAH32226, 437 a.a. ~ 536 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions PDE1B Antibody (monoclonal) (M01) is for research use only and not for use

in diagnostic or therapeutic procedures.


Background

Cyclic nucleotide phosphodiesterases (PDEs) catalyze hydrolysis of the cyclic nucleotides cAMP and cGMP to the corresponding nucleoside 5-prime-monophosphates. Mammalian PDEs have been classified into several families based on their biochemical properties. Members of the PDE1 family, such as PDE1B, are calmodulin (see MIM 114180)-dependent PDEs (CaM-PDEs) that are stimulated by a calcium-calmodulin complex (Repaske et al., 1992 [PubMed 1326532]).


References

1.Distinct metabolism of cyclic adenosine monophosphate in regulatory and helper CD4(+) T cells.Bazhin AV,

Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (36.63 KDa) .



Recombinant ProteinConcentration(ng/tnl)

Detection limit for recombinant GST tagged PDE1B is approximately 0.03ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.