

PI15 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant PI15.

Catalog # AT3304a

Product Information

Application	WB, IHC, E
Primary Accession	O43692
Other Accession	NM_015886
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 Kappa
Clone Names	3B5
Calculated MW	29065

Additional Information

Gene ID	51050
Other Names	Peptidase inhibitor 15, PI-15, 25 kDa trypsin inhibitor, p25TI, Cysteine-rich secretory protein 8, CRISP-8, SugarCrisp, PI15, CRISP8, P25TI
Target/Specificity	PI15 (NP_056970, 18 a.a. ~ 117 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	PI15 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

Background


This gene encodes a trypsin inhibitor. The protein shares similarity to insect venom allergens, mammalian testis-specific proteins and plant pathogenesis-related proteins. It is frequently expressed in human neuroblastoma and glioblastoma cell lines, and thus may play a role in the central nervous system.

References


A genome-wide association study identifies protein quantitative trait loci (pQTLs). Melzer D, et al. PLoS Genet, 2008 May 9. PMID 18464913. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. Generation and initial analysis of more than 15,000

full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932. A novel developmentally regulated gene in lung mesenchyme: homology to a tumor-derived trypsin inhibitor. Kaplan F, et al. Am J Physiol, 1999 Jun. PMID 10362728. cDNA cloning of a novel trypsin inhibitor with similarity to pathogenesis-related proteins, and its frequent expression in human brain cancer cells. Yamakawa T, et al. Biochim Biophys Acta, 1998 Jan 21. PMID 9473672.


Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 kDa).



Immunoperoxidase of monoclonal antibody to PI15 on formalin-fixed paraffin-embedded human kidney. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged PI15 is approximately 0.1 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.