

## PLEK Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant PLEK.

Catalog # AT3340a

### Product Information

---

<b>Application</b>	WB, IF, E
<b>Primary Accession</b>	<a href="#">P08567</a>
<b>Other Accession</b>	<a href="#">BC018549</a>
<b>Reactivity</b>	Human
<b>Host</b>	mouse
<b>Clonality</b>	monoclonal
<b>Isotype</b>	IgG2a Kappa
<b>Clone Names</b>	2D8
<b>Calculated MW</b>	40125

### Additional Information

---


<b>Gene ID</b>	5341
<b>Other Names</b>	Pleckstrin, Platelet 47 kDa protein, p47, PLEK, P47
<b>Target/Specificity</b>	PLEK (AAH18549, 121 a.a. ~ 230 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
<b>Dilution</b>	WB~~1:500~1000 IF~~1:50~200 E~~N/A
<b>Format</b>	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
<b>Storage</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Precautions</b>	PLEK Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.

### References


---

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Multiple common variants for celiac disease influencing immune gene expression. Dubois PC, et al. Nat Genet, 2010 Apr. PMID 20190752. Novel role of pleckstrin homology domain of the Bcr-Abl protein: analysis of protein-protein and protein-lipid interactions. Miroshnychenko D, et al. Exp Cell Res, 2010 Feb 15. PMID 19944685. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. Proteomic identification of pleckstrin-associated proteins in platelets: possible interactions with actin. Baig A, et al. Proteomics, 2009 Sep. PMID 19722192.


## Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .



PLEK monoclonal antibody (M03), clone 2D8. Western Blot analysis of PLEK expression in HL-60 ( (Cat # AT3340a )



Immunofluorescence of monoclonal antibody to PLEK on HeLa cell . [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged PLEK is approximately 3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.