

# PRPF19 Antibody (monoclonal) (M07)

Mouse monoclonal antibody raised against a partial recombinant PRPF19. Catalog # AT3441a

## **Product Information**

Application	WB, IF, E
Primary Accession	<u>Q9UMS4</u>
Other Accession	<u>NM_014502</u>
Reactivity	Human, Mouse, Rat
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	2E6
Calculated MW	55181

### **Additional Information**

Gene ID	27339
Other Names	Pre-mRNA-processing factor 19, 632-, Nuclear matrix protein 200, PRP19/PSO4 homolog, hPso4, Senescence evasion factor, PRPF19 ( <u>HGNC:17896</u> )
Target/Specificity	PRPF19 (NP_055317, 1 a.a. ~ 90 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IF~~1:50~200 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	PRPF19 Antibody (monoclonal) (M07) is for research use only and not for use in diagnostic or therapeutic procedures.

#### Background

PSO4 is the human homolog of yeast Pso4, a gene essential for cell survival and DNA repair (Beck et al., 2008 [PubMed 18263876]).

## References

Regulation of Metnase's TIR binding activity by its binding partner, Pso4. Beck BD, et al. Arch Biochem Biophys, 2010 Jun 15. PMID 20416268.Molecular architecture of the human Prp19/CDC5L complex. Grote M, et al. Mol Cell Biol, 2010 May. PMID 20176811.Blom7alpha is a novel heterogeneous nuclear ribonucleoprotein K homology domain protein involved in pre-mRNA splicing that interacts with SNEVPrp19-Pso4. Grillari J, et al. J Biol Chem, 2009 Oct 16. PMID 19641227.Human Pso4 is a metnase (SETMAR)-binding partner that regulates metnase function in DNA repair. Beck BD, et al. J Biol Chem, 2008 Apr 4. PMID 18263876.Large-scale mapping of human protein-protein interactions by mass spectrometry. Ewing RM, et al. Mol Syst Biol, 2007. PMID 17353931.

#### Images





Western Blot analysis of PRPF19 expression in transfected 293T cell line by PRPF19 monoclonal antibody (M07), clone 2E5.

Lane 1: PRPF19 transfected lysate(55.2 KDa). Lane 2: Non-transfected lysate.



Immunofluorescence of monoclonal antibody to PRPF19 on HeLa cell . [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged PRPF19 is 0.3 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.