

## RAP2B Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant RAP2B.

Catalog # AT3564a

### Product Information

---

<b>Application</b>	WB, IHC, IF
<b>Primary Accession</b>	<a href="#">P61225</a>
<b>Other Accession</b>	<a href="#">BC012362</a>
<b>Reactivity</b>	Human
<b>Host</b>	mouse
<b>Clonality</b>	monoclonal
<b>Isotype</b>	IgG2b kappa
<b>Clone Names</b>	4F12-3C6
<b>Calculated MW</b>	20504

### Additional Information

---

<b>Gene ID</b>	5912
<b>Other Names</b>	Ras-related protein Rap-2b, RAP2B
<b>Target/Specificity</b>	RAP2B (AAH12362, 1 a.a. ~ 183 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
<b>Dilution</b>	WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200
<b>Format</b>	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
<b>Storage</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Precautions</b>	RAP2B Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

### Background

---

This intronless gene belongs to a family of RAS-related genes. The proteins encoded by these genes share approximately 50% amino acid identity with the classical RAS proteins and have numerous structural features in common. The most striking difference between the RAP and RAS proteins resides in their 61st amino acid: glutamine in RAS is replaced by threonine in RAP proteins. Evidence suggests that this protein may be polyisoprenylated and palmitoylated.


### References

---


Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes

Care, 2010 Jul 13. PMID 20628086. Proteome analysis of the thalamus and cerebrospinal fluid reveals glycolysis dysfunction and potential biomarkers candidates for schizophrenia. Martins-de-Souza D, et al. J Psychiatr Res, 2010 May 14. PMID 20471030. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. Targeting of the small GTPase Rap2b, but not Rap1b, to lipid rafts is promoted by palmitoylation at Cys176 and Cys177 and is required for efficient protein activation in human platelets. Canobbio I, et al. Cell Signal, 2008 Sep. PMID 18582561. Identification of genes differentially expressed in human primary lung squamous cell carcinoma. Liu Y, et al. Lung Cancer, 2007 Jun. PMID 17316888.


## Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (45.87 KDa) .



RAP2B monoclonal antibody (M01), clone 4F12-3C6 Western Blot analysis of RAP2B expression in A-431 ( Cat # AT3564a )



Immunoperoxidase of monoclonal antibody to RAP2B on formalin-fixed paraffin-embedded human malignant lymphoma, diffuse large B tissue. [antibody concentration 1 ug/ml]



Immunofluorescence of monoclonal antibody to RAP2B on HeLa cell. [antibody concentration 10 ug/ml]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.